

Original Scientific Paper

New records and noteworthy data of plants, algae and fungi in SE Europe and adjacent regions, 2

Gordana TOMOVIĆ^{1*}, Marko S. SABOVLJEVIĆ^{1*}, Ivana DJOKIĆ¹, Predrag PETROVIĆ², Vladan DJORDJEVIĆ¹, Predrag LAZAREVIĆ¹, Ermin MAŠIĆ³, Senka BARUDANOVIC⁴, Sorin ŠTEFĂNUȚ⁵, Marjan NIKETIĆ⁶, Branislava BUTORAC⁷, Jovana PANTOVIĆ¹, Alma HAJRUDINOVIC-BOGUNIĆ⁸, Faruk BOGUNIĆ⁸, Eva KABAŠ¹, Snežana VUKOJIČIĆ¹, Nevena KUZMANOVIC¹, Sanja Z. DJUROVIĆ⁹ and Uroš BUZUROVIĆ⁶

1 Institute of Botany and Botanical Garden, Faculty of Biology, University of Belgrade, Takovska 43, 11000 Belgrade, Serbia

2 Innovation Centre, Faculty of Technology and Metallurgy, University of Belgrade, Karnegijeva 4, 11000 Belgrade, Serbia

3 Faculty of Science, University of Sarajevo, Zmaja od Bosne 33-35, 71000 Sarajevo, Bosnia and Herzegovina

4 Centre for Ecology and Natural Resources "Academician Sulejman Redžić", Zmaja od Bosne 33-35, 71000 Sarajevo, Bosnia and Herzegovina

5 Bucharest Institute of Biology, Romanian Academy, 296 Splaiul Independentei, 060031 Bucharest, P.O. Box 56-53, Romania

6 Natural History Museum, Njegoševa 51, 11000 Belgrade, Serbia

7 Institute for Nature Conservation of the Vojvodina Province, 21000 Novi Sad, Serbia

8 Faculty of Forestry, University of Sarajevo, Zagrebačka 20, 71000 Sarajevo, Bosnia and Herzegovina

9 Faculty of Agriculture, University of Niš, Kosančićeva 4, 37000 Kruševac, Serbia

* column editors, to whom contributions should be sent (botanicaserbica@bio.bg.ac.rs)

ABSTRACT:

This paper presents new records and noteworthy data on the following taxa in SE Europe and adjacent regions: the diatom alga *Eunotia boreoalpina*; the saprotrophic fungus *Clitocybe truncicola*; the liverwort *Haplomitrium hookeri*; the moss *Leptodon smithii*; the monocots *Epipactis purpurata*, *Stipa tirsa*, *Typha laxmannii* and *T. shuttleworthii*; and the dicots *Krascheninnikovia ceratoides*, *Polygonum albanicum* and *Sorbus latifolia*.

Keywords:

new record, *Clitocybe truncicola*, *Eunotia boreoalpina*, *Epipactis purpurata*, *Haplomitrium hookeri*, *Krascheninnikovia ceratoides*, *Leptodon smithii*, *Polygonum albanicum*, *Sorbus latifolia*, *Stipa tirsa*, *Typha laxmannii*, *Typha shuttleworthii*

UDC: 581.95: 582.261.1+582.28+582.3/99

Revision accepted: 04 September 2020

***Clitocybe truncicola* (Peck) Sacc., fam. Tricholomataceae (fungus, saprotrophic)**

Contributors: Ivana DJOKIĆ and Predrag PETROVIĆ

Geographical focus: Serbia

New record and noteworthy data: The first record of *Clitocybe truncicola* in Serbia.

Specimen data: Belgrade area, Ada Ciganlija, N 44.782283°, E 20.379246°, on a decaying log of a decidu-

ous tree, in the protected area "Fungi of Ada Ciganlija", 84 m a.s.l., 11 October 2019; leg. Djokić I.; det. Petrović P.

Voucher: Natural History Museum in Belgrade, National Fungal Collection (BEO) 21128.

From early October to the end of November 2019, trunk funnel (*C. truncicola*) basidiocarps were collected from several sites on Ada Ciganlija, a river island

that has been artificially turned into a peninsula, located in the Sava River's course through central Belgrade. Specifically, basidiocarps were recorded within a protected area of 21 ha, a central forested complex declared the "Fungi of Ada Ciganlija" protected habitat in 2013. This area is specific as it hosts more than 250 species of macromycetes, many of which are included on Serbian and international lists of rare or endangered species. *Clitocybe truncicola* was originally described from North America (BAS *et al.* 1995) and the first finds of this species in Europe were reported as recently as in 1988 (BREITENBACH & KRANZLIN 1992). Since then it was reported from several European countries (Austria, Czech Republic, Denmark, Estonia, France, Germany, Hungary, Italy, Spain, the Netherlands, the United Kingdom), in some being considered very rare. It is one of only two European *Clitocybe* species, alongside *C. subbulbipes* Murr., that occur on dead wood of various tree species from the genera *Acer*, *Alnus*, *Carpinus*, *Fraxinus*, *Populus*, and *Ulmus*, as well as on *Sorbus aucuparia* L., which is why it was confused in the past with the macroscopically and ecologically similar *Ossicaulis lignatilis* (Pers.) Redhead & Ginns [formerly known as *Clitocybe lignatilis* (Pers.) P. Karst.] , also found in the same habitat on Ada Ciganlija (BAS *et al.* 1995; LOHMEYER 1999). According to the Global Biodiversity Information Facility (GBIF database – <https://www.gbif.org/>), *C. truncicola* has two disjunct records, one in China (reported in 2014) and another from Morocco (reported in 1989). The new record of this species is important both as a new finding for the protected area on Ada Ciganlija, an area covered by active measures on fungal conservation, and as a first record for Serbia.

***Epipactis purpurata* Sm., fam. Orchidaceae (monocot, vascular plant)**

Contributors: Vladan DJORDJEVIĆ and Predrag LAZAREVIĆ

Geographical focus: Serbia

New records and noteworthy data: These are the first records for the Ivanjica region and Mt. Zlatibor, at the same time representing the first findings of this species within the MGRS 34T DP2 50 × 50 km² UTM grid cell. The new locality in the village of Lisa near the town of Ivanjica represents the easternmost limit of the species' distribution in the Dinaric part of the Balkan Peninsula.

Specimen data: 1) western Serbia, Ivanjica, village of Lisa, Samograd – Straža, N 43.661703°, E 20.231065°, MGRS 34T DP33, *Fagetum montanum*, limestone, exp. NW, incl. 20°, 810 m a.s.l., 21 July 2019; leg./det. Djordjević V. 17684; 2) western Serbia, Mt. Zlatibor, Čigota peak, Brezanje, N 43.65965°, E 19.77522°, MGRS 34T DP03, *Fagetum montanum*, limestone, exp. NE, incl. 20°, 1035 m, m a.s.l., 13 July 2020; leg. Djordjević V.; det. Djordjević V., Lazarević P. 17685; 3) western Serbia, Mt. Zlatibor,

Murtenica peak, Šišačka Kosa, N 43.60519°, E 19.75772°, MGRS 34T CP92, *Fagetum montanum*, limestone, exp. S, incl. 25°, 1220 m a.s.l., 14 July 2020; leg. Djordjević V.; det. Djordjević V., Lazarević P. 17686; 4) western Serbia, Mt. Zlatibor, Murtenica peak, Brijač, Velika Stojkovača, N 43.59115°, E 19.78354°, MGRS 34T DP02, *Abieti-Fagetum*, limestone, exp. NE, incl. 25°, 1290 m a.s.l., 14 July 2020; leg. Djordjević V.; det. Djordjević V., Lazarević P. 17687; 5) western Serbia, Mt. Zlatibor, Sirogojno (church), N 43.68913°, E 19.88140°, MGRS 34T DP03, *Fagetum montanum*, limestone, exp. NE, incl. 25°, 865 m a.s.l., 22 August 2020; leg./det. Djordjević V. 17688.

Vouchers: Herbarium of the Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade (BEOU), vascular plant collection 17684, 17685, 17686, 17687, 17688.

This species is distributed primarily in the temperate zone of Western and Central Europe; it occurs north to Denmark, Great Britain and the Baltic States, west to France and northern Spain, and south to southern Italy, Greece and Bulgaria (RANKOU 2011). The species was found on Mt. Maljen (Bukovi) for the first time in Serbia (DJORDJEVIĆ *et al.* 2010), and later on Mts. Jablanik, Tara, Zvijezda, Pobijenik, Jadovnik and Kamena Gora (DJORDJEVIĆ *et al.* 2017). New findings of this species in the Ivanjica region and on Mt. Zlatibor represent four new 10 × 10 km UTM grid cells in the region of western Serbia, in addition to previous findings only on Mts. Tara and Zvijezda (DJORDJEVIĆ *et al.* 2017).

The population in Ivanjica (village of Lisa) numbered two individuals within an area of 20 m², whereas the populations on Mt. Zlatibor consisted of the following numbers of individuals: four individuals / 20 m² (Murtenica peak, Šišačka kosa); seven individuals / 30 m² (Murtenica peak, Brijač); eight individuals / 50 m² (Sirogojno); and 16 individuals / 200 m² (Čigota, Brezanje). The species is endangered (IUCN: EN) in Bulgaria, the Czech Republic and Liechtenstein, whereas it has the status of a vulnerable species (IUCN: VU) in Slovakia, Luxembourg and Austria (KULL *et al.* 2016). According to DJORDJEVIĆ *et al.* (2017), the estimated IUCN conservation status of this species in Serbia is vulnerable (VU). Accordingly, the species should be included in the Red Data Book of Flora of Serbia.

***Eunotia boreoalpina* Lange-Bertalot et Nörpel-Schempp in Lange-Bertalot & Metzeltin, fam. Eu-notiaceae (diatom, algae)**

Contributors: Ermin MAŠIĆ and Senka BARUDANOVIĆ
Geographical focus: Bosnia and Herzegovina

New record and noteworthy data: First record for Bosnia and Herzegovina. Species recorded in peatlands on Mt. Vranica.

Specimen data: Mt. Vranica, species recorded in four small mountain peat bogs: 1) N 43.95192°, E 17.75827°,

1714 m a.s.l.; **2)** N 43.95134°, E 17.76057°, 1755 m a.s.l.; **3)** N 43.95131° E 17.76065°, 1755 m a.s.l.; **4)** N 43.95137°, E 17.76060°, 1761 m a.s.l.), August and September 2019; leg./det. Mašić E.

Voucher: Diatom collection (Mašić E.) s/n, Laboratory for study of the systematics of algae and fungi, Department of Biology, Faculty of Science, University of Sarajevo (Bosnia and Herzegovina).

The first systematic algological research on Mt. Vranica (Bosnia and Herzegovina) was carried out in different wet habitats developed near mountain springs and brooks (KAPETANOVIĆ & HAFNER 2007). Additional data on the diversity of algae in freshwater habitats on Mt. Vranica were published by various authors (BARUDANOVIĆ et al. 2015; MAŠIĆ et al. 2019).

The presence of the species *Eunotia boreoalpina* in freshwater habitats on Mt. Vranica was expected. According to published data, this species in Central Europe is scattered but locally abundant in mountainous regions on siliceous bedrock, in moderately acidic electrolyte-poor springs, headwaters, lakes and minerotrophic fens (CANTONATI et al. 2017).

In Bosnia and Herzegovina (Mt. Vranica), the species *E. boreoalpina* was identified only from mountain peat bogs. According to KRIZMANIĆ et al. (2015), this species is characteristic of moderately acidic waters with low electrolyte content. Altitudes where *E. boreoalpina* was identified range from 1714 to 1761 m a.s.l. In peat bogs where the species was identified, pH ranged from 4.16 to 5.93, dissolved oxygen ranged from 7.00 to 7.82 mg·L⁻¹, while specific conductivity ranged from 22.44 to 600.8 µS·cm⁻¹. According to the German Red List (LANGE-BERTALOT & STEINDORF 1996), this species is not classified as threatened, but in Bulgaria (within the Balkan Peninsula) STOYNEVA-GÄRTNER (2015) assessed its status as vulnerable (VU). Unfortunately, the peat bogs where *E. boreoalpina* was identified are today under strong anthropogenic influences. These habitat types cover only a small part of Mt. Vranica, and as they are the most fragile and the most sensitive habitat in this area, they deserve special conservation attention.

Haplomitrium hookeri (Sm.) Nees, fam. Haplomitriaceae (liverwort, bryophyte)

Contributor: Sorin ȘTEFĂNUȚ

Geographical focus: Romania

New record and noteworthy data: This is the first record for the Parâng Mountains.

Specimen data: Southern Carpathians, Mts. Parâng, on a bank of Lung Lake, N 45.34558333°, E 23.55155556°, 2023 m a.s.l., 28 June 2012; leg./det. Ștefănuț S.

Voucher: Herbarium of the Bucharest Institute of Biology, Romanian Academy (BUCA), bryophyte collection, B4371.

The first published record of *Haplomitrium hookeri* for Romania was from the Rodna Mountains (ȘTEFĂNUȚ 2000), and since then three new reports were published: on the Bucegi Massif (ȘTEFĂNUȚ 2007), and on Mts. Făgăraș and Retezat (ȘTEFĂNUȚ 2010).

The presence of *H. hookeri* in the Parâng Mountains was expected, thus completing the species' distribution in all of the Southern Carpathians. This is the southernmost record of the given species in Southeast Europe.

According to the European Red List of Mosses, Liverworts and Hornworts (HODGETTS et al. 2019), *H. hookeri* has been assessed as being of least concern (LC), but for Romania it is assigned to the category of vulnerable (VU) (ȘTEFĂNUȚ & GOIA 2012).

Krascheninnikovia ceratoides (L.) Gueldenst., fam. Amaranthaceae (dicot, vascular plant)

Contributors: Marjan NIKETIĆ and Branislava BUTORAC

Geographical focus: Serbia

New records and noteworthy data: On the basis of extensive field investigation and herbarium revision, the species should be excluded from the flora of Serbia.

Specimen data: Based on revision of the specimen of the species reported in Serbia and checking the reported locality, this species cannot be confirmed for Serbia.

Vouchers: The previous voucher specimen [Banat, village of Banatska Palanka, Danube-Tisza-Danube Canal, MGRS 34T EQ26, 15 September 1978; leg./det. Joksimović Ž. (sub *Eurotia ceratoides*) 100003 (Natural History Museum in Belgrade, General Herbarium of the Balkan Peninsula – BEO)] was revised and found to actually belong to the naturalised species *Bassia scoparia* (L.) A. J. Scott (rev. Niketić M.).

The only record of the presence of Pamirian winterfat (*K. ceratoides*, tribe Axyrideae) in Serbia originates from JOKSIMOVIĆ (1977), and it was subsequently cited in Flora of SR Serbia (DIKLJ & NIKOLIĆ 1986), but its occurrence has not been confirmed to date. An inspection of the original herbarium material in BEO showed that it was based on one specimen in a piece of newsprint, on the margin of which was written "*Eurotia ceratoides*". However, this specimen was misidentified and actually belongs to *Bassia scoparia* (L.) A. J. Scott. Therefore, neither Joksimović's record nor the data from the publication Flora of SR Serbia can be taken as valid, and the name *K. ceratoides* should be treated as misapplied in floristic data for Serbia. This Holarctic species is very rare in the European flora (JALAS & SUOMINEN 1980) and survived in remains of the Pleistocene cold-steppe and semi-desert flora (NIKETIĆ 2018). According to BARTHA & KIRÁLY (2015), it became extinct in the Hungarian flora, and thus in that of the Pannonian Plain. The only enclave on the Balkan Peninsula is located in the central part of North Macedonia (JALAS & SUOMINEN 1980).

***Leptodon smithii* (Hedw.) Web. & Mohr, fam. Neckeraceae (moss, bryophyte)**

Contributors: Jovana PANTOVIĆ and Marko SABOVLJEVIĆ

Geographical focus: Serbia

New and noteworthy data: Two recent records for Serbia, the first record for Vojvodina province (North Bačka county, Bačka region) and the first one for the region of western Serbia (Zlatibor county).

Specimen data: 1) Subotičko Horgoška sands, Bukvač, N 46.157408°, E 19.686411°, on *Fraxinus* bark in a mixed deciduous forest, 130 m a.s.l., 21 May 2020; leg. Pantović J., Stevanoski I., Bogosavljević J.; det. Pantović J.; rev. Sabovljević M. 07875; 2) Tara National Park, gorge of the Grlac River, N 43.97778°, E 19.33819°, on a fallen trunk in a canyon with mixed thermophilous vegetation, 406 m a.s.l., 11 July 2016; leg./det. Pantović J.; rev. Sabovljević M. 07876.

Voucher: Herbarium of the Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade Herbarium (BEOU), bryophyte collection, Bryo 07875 and 07876.

Leptodon smithii is distinctive epiphytic moss that grows on the trunks or larger branches of mature trees in woods, or more rarely on basic rocks (ATHERTON *et al.* 2010). This Mediterranean-Atlantic species is notably tolerant of dry shade, and it is common in the Mediterranean region (BLOCKEEL *et al.* 2014). On the European level, it is considered to be of least concern (LC) according to the new European Red List of Mosses, Liverworts and Hornworts (HODGETTS *et al.* 2019). However, it is regionally rare and protected in some of Serbia's neighbours, e.g., it is assessed as CR in Romania (ȘTEFĂNUȚ & GOIA 2012) and EN in Hungary (PAPP *et al.* 2010).

This species was previously known in Serbia, but from only two sites. The first one, and the only record for the central part of Serbia, was given more than 70 years ago by Soška (SOŠKA 1949), who reported it from the Topčider Park in Belgrade. However, this species was not refound in recently conducted exploration of the Belgrade metropolitan area (SABOVLJEVIĆ & GRDOVIĆ 2009), and the Soška herbarium seems to be ruined. The second historical locality is a spring of the river Istočka Reka, in Kosovo and Metohija. This record was published by MARTINČIĆ (2006) on the basis of more than 50-year-old herbarium samples. The two recent records are a significant contribution to the bryophyte flora of Serbia and to knowledge of the distribution of bryophytes in Serbia, as well as possible bio-documentation of climate change and global warming.

***Polygonum albanicum* Jav., fam. Polygonaceae (dicot, vascular plant)**

Contributors: Marjan NIKETIĆ and Gordana TOMOVIĆ

Geographical focus: Serbia

New records and noteworthy data: These are the first records for Serbia proper and several new localities for the Republic of Serbia.

Specimen data: 1) western Serbia, Mt. Zlatibor, Partizanske Vode – Semegnjevo, MGRS 34T CP94, serpentinite, 16 July 1989; leg./det. Niketić M. s/n. (BEO); 2) southwestern Serbia, Mt. Ozren, Goleč peak, N 43.2531869°, E 19.9136436°, MGRS 34T DN18, tiny screes by the roadside, harzburgite, 1325 m a.s.l., 20 July 2020; leg./det. Niketić M., Tomović G. 68407 (BEOU), s.n. (BEO); 3) southwestern Serbia, Mt. Ozren, Orlujak peak, N 43.2522335°, E 19.8976081°, MGRS 34T DN18, tiny screes by the roadside, harzburgite, 1330 m a.s.l., 20 July 2020; leg./det. Niketić M., Tomović G. 68408 (BEOU), s.n. (BEO); 4) southwestern Serbia, Mt. Ozren, Pape peak, N 43.2382985°, E 19.9059896°, MGRS 34T DN18, tiny screes by the roadside, harzburgite, 1345 m a.s.l., 20 July 2020; leg./det. Niketić M., Tomović G. s/n. (BEO); 5) southwestern Serbia, Mt. Ozren, Revuša peak, N 43.2448967°, E 19.8535345°, MGRS 34T DN08, tiny screes by the roadside, harzburgite, 1630 m a.s.l., 21 July 2020; leg./det. Niketić M., Tomović G. 68409 (BEOU), s/n. (BEO)

Vouchers: Herbarium of the Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade (BEOU), vascular plant collection 68407, 68408, 68409; Natural History Museum in Belgrade, General Herbarium of the Balkan Peninsula (BEO), s/n.

The presence of this species in Serbia was first recorded by JÁVORKA *et al.* (1926) from the Kosovo & Metohija province (Djakovica, vicinity of the village of Morina, Čafa Morina pass). After JÁVORKA *et al.* (1926), it was confirmed for the same locality by WRABER (1990). Until now, this species was not known for Serbia proper, and these five records represent the first reports of its presence in this area. At the same time, the newly-discovered locality of *P. albanicum* on Mt. Zlatibor represents the northernmost and most continental border of the species' distribution.

This obligate serpentinophyte inhabits rocky grounds and tiny screes by roadsides in hilly regions. On Mt. Zlatibor and at all four localities on Mt. Ozren, the sub-populations are very poor and consist of only a few flowering individuals.

***Sorbus latifolia* (Lam.) Pers. *sensu lato* (*S. aria* s.l. × *S. torminalis*), fam. Rosaceae (dicot, vascular plant)**

Contributors: Alma HAJRUDINOVIĆ-BOGUNIĆ and Faruk BOGUNIĆ

Geographical focus: Montenegro and Serbia

New records and noteworthy data: This is the first record for Montenegro and the second one reported for Serbia.

Specimen data: 1) Northwestern Montenegro, Nikšić, Grahovac, N 42.6969444°, E 18.6297222°, 1010 m a.s.l.,

5 October 2019; leg./det. Hajrudinović-Bogunić A., Bogunić F. 60020; $2n = 3x$; 2) eastern Serbia, Mt. Suva Planina, Devojački Grob, N 43.1988889°, E 22.1416667°, 1300 m a.s.l., 4 October 2018.; leg. Bogunić F.; det. Bogunić F., Hajrudinović-Bogunić A. 60021; $2n = 3x$;

Vouchers: National Museum of Bosnia and Herzegovina, Herbarium of Bosnia and Herzegovina (SARA), the Herbarium Europaeum collection 60020, 60021.

Despite extensive investigation of the Balkan flora, gaps in taxonomic and distribution knowledge are still evident in certain plant groups, particularly in taxonomically complicated ones (ŠPANIEL *et al.* 2017). The genus *Sorbus* is a typical example, with a complex and unresolved taxonomy which is far from a clear-cut situation based on solid knowledge in the Balkans. The exceptional genetic and morphological diversity within the genus is a result of recurrent polyploidisation, homo- and heteroploid hybridisation, backcrosses and apomixis (ROBERTSON *et al.* 2010; HAJRUDINOVIC *et al.* 2015a), which encouraged description of novel taxa across the European continent (LEPŠÍ *et al.* 2008, 2009; HAJRUDINOVIC *et al.* 2015b; NÉMETH *et al.* 2016).

Sorbus subgenus *Tormaria* Májovský & Bernátová represents a complex aggregate containing taxa which originated through spontaneous hybridisation and backcrosses between members of the subgenera *Aria* and *Torminaria* (RICH *et al.* 2010). Central and Western Europe are considered as hotspots of *Tormaria* diversity that include various, mostly polyploid microspecies under the common name *S. latifolia* (Lam) Pers. s.l. (KURTO 2009). However, the distribution of *Tormaria* members is much wider and encompasses the Balkans as well (HAJRUDINOVIC *et al.* 2012; ZIELIŃSKI & VLADIMIROV 2013; TEOFILOVSKI 2017).

Recently, two new localities of members of the subgenus *Tormaria* on the Balkan Peninsula were discovered in Serbia and Montenegro. Analysis of nuclear microsatellites of individuals from both localities confirmed triploids (HAJRUDINOVIC-BOGUNIĆ, preliminary results). These findings represent the first known locality for Montenegro and the second one reported for Serbia. The first locality of *S. latifolia* s.l. in Serbia, based on taxonomic revision of a single herbarium voucher, was reported by ZIELIŃSKI & VLADIMIROV (2013).

Stipa tirsa Steven, fam. Poaceae (monocot, vascular plant)

Contributors: Eva KABAŠ and Snežana VUKOJIČIĆ

Geographical focus: Serbia

New record and noteworthy data: This is the first record for the region of central Serbia.

Specimen data: central Serbia, Mt. Kopaonik, Krmeljica peak, N 43.262793°, E 20.741679°, 1016 m a.s.l., 27 May 2015; leg./det. Kabaš E.; rev. Vukojičić S. 41977.

Voucher: Herbarium of the Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade (BEOU), vascular plant collection 41977.

According to MARTINOVSKY (1980), *Stipa tirsa* is present in the Balkans only at certain localities, representing the western borders of its distribution. Until now, *S. tirsa* was known only from a few localities in Serbia – Mt. Rtanj, Mt. Suva Planina and the spa Brestovačka Banja in eastern Serbia (VUKIĆEVIĆ 1976), as well as the Mt. Vršačke Planine in the province of Vojvodina, where it was found in meadow steppe communities (VUČKOVIĆ 1991).

The newly recorded population on Mt. Kopaonik numbers up to 50 individuals and is found at one small site on steep slopes of the Krmeljica peak on an ultramafic substrate within a dry grassland steppe-like stand of *Stipa mayeri* Martinovsky. Because *S. tirsa* is a steppe relic species with very narrow distribution limited to restricted patches, the sites of its occurrence are of great conservation importance.

Typha laxmannii Lepech., fam. Typhaceae (monocot, vascular plant)

Contributor: Nevena KUZMANOVIĆ

Geographical focus: Serbia

New records and noteworthy data: This is the first record for the region of western Serbia. Neophyte, probably spreading out.

Specimen data: western Serbia, Zlatibor county, Mt. Zlatibor, Bregovi, N 43.703966°, E 19.689724°, shallow pond beside the main road (European route E763), 1008 m a.s.l., 22 August 2020; leg./det. Kuzmanović N. 54494.

Voucher: Herbarium of the Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade (BEOU), vascular plant collection 54494.

The first records of *Typha laxmannii* for the Serbian flora were reported by BUDAK (1986) for the territory of Vojvodina province, where many localities in the Banat and Bačka regions were listed. More recent findings include records from eastern and southeastern Serbia: the village of Šaprance, environs of Dimitrovgrad, Trgovište – Vražji Kamen, environs of Vranjska Banja (ZLATKOVIĆ *et al.* 2007), as well as northeastern Serbia: the town of Zaječar – village of Rgotina, Rgotsko Lake (ZLATKOVIĆ & BOGOŠAVLJEVIĆ 2014). Records of *T. laxmannii* are lacking from western Serbia. The first record for western Serbia (from Zlatibor county) is given here. The species was found in a shallow pond beside the main road.

Typha shuttleworthii Koch & Sonder, fam. Typhaceae (monocot, vascular plant)

Contributors: Sanja Z. DJUROVIĆ and Uroš BUZUROVIĆ

Geographical focus: Serbia

New records and noteworthy data: New sites in central and western Serbia are given for *T. shuttleworthii*, which

is considered to be endangered in Serbia. It is a new species in the Tara National Park, the Golija Nature Park and the Radan Nature Park. The species is listed in the Bern Convention.

Specimen data: 1) western Serbia, Mt. Tara, Kurjačine, MGRS 34T CP66, 5 August 2017; leg. Vukojičić S., Kabaš E., Djurović S., Buzurović U.; det. Đurović, S. 52007 (BEOU); 2) western Serbia, Mt. Tara, Dikava – Kozulja, N 43.89517°, E 19.323501°, MGRS 34T CP66, 1289 m a.s.l., 25 July 2019; leg./det. Djurović S., Buzurović U. 67650 (BEOU), 85003 (BEO); 3) western Serbia, Mt. Golija, Bzovik, Izubra, N 43.420927°, E 20.431155°, MGRS 34T DP50, 1102.5 m a.s.l., 16 July 2019; leg./det. Vukojičić S., Kabaš E., Veljić M., Lazarević P., Djurović S. 67820 (BEOU); 4) central Serbia, Mt. Radan, Gornji Gajtan, N 42.9815409°, E. 21.4720178°, MGRS 34T EN35, 1036 m a.s.l., 30 July 2020; leg./det. Djurović S., Buzurović U., Veljić M. 68671 (BEOU), 85002 (BEO); 5) central Serbia, Mt. Radan, Kosmača, N 43.0933059°, E. 21.375447°, MGRS 34T EN37, 800 m a.s.l., 31 July 2020; Djurović S., Buzurović U. (*photo documentation*).

Vouchers: Herbarium of the Institute of Botany and Botanical Garden "Jevremovac", University of Belgrade (BEOU), vascular plant collection 52007, 67650, 67820, 68671; Natural History Museum in Belgrade, General Herbarium of the Balkan Peninsula (BEO) 85003, 85002; photo documentation: S. Djurović, U. Buzurović.

Typha shuttleworthii is covered by the Bern Convention and included on national red lists and/or in red data books in many European countries (KORNECK *et al.* 1996; KÄSERMANN 1999; KIRÁLY 2007; HLAVÁČEK & GRULICH 2009; FELBABA-KLUSHYNA 2011; KOZŁOWSKA *et al.* 2011). In European Red List of Vascular Plants Needing Protection, it is marked as "Data Deficient" (BILZ *et al.* 2011). In Serbia it was first considered critically endangered with only four recorded sites: the environs of Novi Sad, Kragujevac and Sjenica; and Mt. Kukavica (RANDJELOVIĆ 1999 and references therein). With emerging new sites in eastern Serbia (Crna Reka gorge, Svrliški Timok gorge and Topli Dol), the threat status of this species in Serbia was changed to endangered (EN; TOMOVIĆ *et al.* 2009 and references therein).

Although four new sites are recorded, the species forms relatively small patches, usually not covering more than a few square metres, mostly in roadside ditches (except on Mt. Golija, where it also grows along a stream). The species is probably often not distinguished from *T. latifolia* L., and further localities will no doubt be discovered in Serbia.

REFERENCES

- ATHERTON I, BOSANQUET SD & LAWLEY M (eds.). 2010. *Mosses and liverworts of Britain and Ireland: a field guide*. British Bryological Society, Plymouth.
- BARTHA D & KIRÁLY G (eds.). 2015. *Distribution atlas of vascular plants of Hungary*. University of West Hungary Press, Sopron.
- BARUDANOVIĆ S, MAŠIĆ E & MACANOVIĆ A. 2015. *Trešetišta na bosanskim planinama*. Prirodno-matematički fakultet, Univerzitet u Sarajevu.
- BAS C, KUYPER T, NOORDELOOS M & VELLINGA E. 1995. *Flora Agaricina Neerlandica* 3. Balkema, Rotterdam.
- BILZ M, KELL SP, MAXTED N & LANSDOWN RV. 2011. *European Red List of Vascular Plants*. Publications Office of the European Union, Luxembourg.
- BLOCKEEL TL, BOSANQUET SD, HILL MO & PRESTON CD. 2014. *Atlas of British & Irish Bryophytes Volume 2*. Pisces publications, Newbury.
- BREITENBACH J & KRANZLIN F. 1992. *Fungi of Switzerland 3: Boletes and agarics 1st part*. Edition Mycologia Lucerne, Lucerne.
- BUDAK V. 1986. *Typha laxmannii* Lepech. In: SARIĆ MR & DIKLIĆ N (eds.), *Flora SR Srbije* 10, pp. 255–256, Srpska akademija nauka i umetnosti, Beograd.
- CANTONATI M, KELLY GM & LANGE-BERTALOT H. 2017. *Freshwater Benthic Diatoms of Central Europe: Over 800 Common Species Used in Ecological Assessment. English edition with updated taxonomy and added species*. Koeltz Botanical Book, Oberreifenberg.
- DIKLIĆ N & NIKOLIĆ V. 1986. *Krascheninnikovia Gueldenst.* In: SARIĆ M & DIKLIĆ N (eds.), *Flora SR Srbije* 10, Dodatak (2), pp. 54–55, Srpska akademija nauka i umetnosti, Beograd.
- DJORDJEVIĆ V, TOMOVIĆ G & LAKUŠIĆ D. 2010. *Epipactis purpurata* Sm. (Orchidaceae) – a new species in the flora of Serbia. *Archives of Biological Science* 62: 1175–1180.
- DJORDJEVIĆ V, LAKUŠIĆ D, JOVANOVIĆ S & STEVANOVIĆ V. 2017. Distribution and conservation status of some rare and threatened orchid taxa in the central Balkans and the southern part of the Pannonian Plain. *Wulfenia* 24: 143–162.
- FELBABA-KLUSHYNA L. 2011. *Typha shuttleworthii* in Ukraine and adjoining regions: tendencies of dynamics of distribution, ecological and coenotic peculiarities. *Botanica Serbica* 35(2): 121–124.
- HAJRUDINOVIC A, BAŠIĆ N & BOGUNIĆ F. 2012. *Sorbus latifolia* (Rosaceae): a new species in the flora of Bosnia and Herzegovina. *Academy of Sciences and Arts of Bosnia and Herzegovina, Special Editions CXLVIII, Proceedings* 22: 175–186.
- HAJRUDINOVIC A, FRAJMAN B, SCHÖNSWETTER P, SILAJDŽIĆ E, SILJAK-YAKOVLEV S & BOGUNIĆ F. 2015b. Towards a better understanding of polyploid *Sorbus* (Rosaceae) from Bosnia and Herzegovina (Balkan Peninsula), including description of a novel, tetraploid apomictic species. *Botanical Journal of the Linnean Society* 178: 670–685.

- HAJRUDINOVIC A, SILJAK-YAKOVLEV S, BROWN SC, PUSTAHIJA F, BOURGE M, BALLIAN D & BOGUNIĆ F. 2015a. When sexual meets apomict: genome size, ploidy level and reproductive mode variation of *Sorbus aria* s.l. and *S. austriaca* (Rosaceae) in Bosnia and Herzegovina. *Annals of Botany* **116**(2): 301–312.
- HLAVÁČEK R & GRULICH V. 2009. *Typha shuttleworthii* W. D. J. Koch & Sond. In: HLADINEC J & LUSTYK P (eds.), Additions to the flora of the Czech Republic. VIII. *Zprávy České Botanické Společnosti* **44**: 310–312.
- HODGETTS N, CALIX M, ENGLEFIELD E, FETTES N, GARCIA CRIADO M, PATIN L, NIETO A, BERGAMINI A, BISANG I, BAISHEVA E, CAMPISI P, COGONI A, HALLINGBACK T, KONSTANTINOVA N, LOCKHART N, SABOVLJEVIC M, SCHNYDER N, SCHROCK C, SERGIOC, SIM SIM M, VRBA J, FERREIRA CC, AFONINA O, BLOCKEEL T, BLOM H, CASPARI S, GABRIEL R, GARCIA C, GARILLETI R, GONZALEZ MANCEBO J, GOLDBERG I, HEDENAS L, HOLYOAK D, HUGONNOT V, HUTTUNEN S, IGNATOV M, IGNATOVA E, INFANTE M, JUUTINEN R, KIEBACHER T, KOCKINGER H, KUČERA J, LONNELL N, LUTH M, MARTINS A, MASLOVSKY O, PAPP B, PORLEY R, ROTHERO G, SODERSTROM L, ŠTEFĀNUŠ S, SYRJANEN K, UNTEREINER A, VAŇA J, VANDERPOORTEN A, VELLAK K, ALEFFI M, BATES J, BELL N, BRUGUES M, CRONBERG N, DENYER J, DUCKETT J, DURING HJ, ENROTH J, FEDOSOV V, FLATBERG KI, GANEVA A, GORSKI P, GUNNARSSON U, HASSEL K, HESPAÑOL H, HILL M, HODD R, HYLANDER K, INGERPUU N, LAAKA-LINDBERG S, LARA F, MAZIMPAKA V, MEŽAKA A, MULLER F, ORGAZ JD, PATINO J, PILKINGTON S, PUCHE F, ROS RM, RUMSEY F, SEGARRA-MORAGUES JG, SENECA A, STEBEL A, VIRTANEN R, WEIBULL H, WILBRAHAM J & ŽARNOWIEC J. 2019. *A miniature world in decline: European Red List of Mosses, Liverworts and Hornworts*. International Union for Conservation of Nature (IUCN), Brussels.
- JALAS J & SUOMINEN J. 1980. *Atlas Flora Europaea. Distribution of vascular plants in Europe 5 (Chenopodiaceae to Basellaceae)*. The Committee for Mapping the Flora of Europe & Societas Biologica Fennica Vanamo, Helsinki.
- JÁVORKA S, DEGEN A, GÁYER Gy, SOÓ R, TRAUTMANN R & ZAHN KH. 1926. Anthophyta [In: CSIKI E, JÁVORKA S & KÜMMERLE JB. *Adatok Albánia flórájához*], In: TELEKI P & CSIKI E. (eds.), *A Magyar tudományos akadémia Balkán-Kutatásainak Tudományos Eredményei* 3, pp. 219–346, A Magyar Tudományos Akadémia Kiadása, Budapest.
- JKOSIMOVIĆ Ž. 1977. *Eurotia ceratooides* (L.) C. A. Mey. – nova biljna vrsta u flori SR Srbije. *Glasnik Prirodnjačkog muzeja u Beogradu* B **32**: 79–80.
- KAPETANOVIĆ T. & HAFNER D. 2007. Diatoms of wet habitats in the subalpine belt of Mt. Vranica (Bosnia and Herzegovina). In: KUSBER WH & JAHN R (eds.), Proceedings of the 1st Central European Diatom Meeting. Botanic Garden and Botanical Museum Berlin-Dahlem, Freie Universität Berlin, pp. 73–78.
- KÄSERMANN C. 1999. VU *Typha shuttleworthii* W. D. J. Koch & Sond. – Shuttleworths Rohrkolben – Typhaceae. Published on the Internet https://www.infoflora.ch/assets/content/documents/merkblaetter_artenschutz_de/typ_ [Accessed 12 August 2020]. shut_d.pdf
- KIRÁLY G. (ed.) 2007. *Red list of the vascular flora of Hungary*. Private edition, Sopron.
- KORNECK D, SCHNITTNER M & VOLLMER I. 1996. Rote Liste der Farn- und Blütenpflanzen (Pteridophyta et Spermatophyta) Deutschlands. *Schriftenreihe für Vegetationskunde* **28**: 21–187.
- KOZŁOWSKA K, NOBIS A & NOBIS M. 2011. *Typha shuttleworthii* (Typhaceae), new for Poland. *Polish Botanical Journal* **56**: 299–305.
- KRIZMANIĆ J, ILIĆ M, VIDAKOVIĆ D, SUBAKOV SIMIĆ G, CVETANOVIĆ K & PETROVIĆ J. 2015. New records and rare taxa of the genus *Eunotia* Ehrenberg (Bacillariophyceae) for the diatom flora of Serbia. *Botanica Serbica* **39**(1): 35–44.
- KULL T, SELGIS U, PECINA MV, METSARE M, ILVES A, TALI K & SHEFFERSON RP. 2016. Factors influencing IUCN threat levels to orchids across Europe on the basis of national red lists. *Ecology and Evolution* **6**: 6245–6265.
- KURTTO A. 2009. Rosaceae (pro parte majore). In: *Euro + Med Plantbase – the information resource for Euro-Mediterranean plant diversity*. Published on the Internet <http://ww2.bgbm.org/EuroPlusMed/> [Accessed 14 July 2020].
- LANGE-BERTALOT H & STEINDORF A. 1996. Rote Liste der limnischen Kieselalgen (Bacillariophyceae) Deutschlands. *Schriftenreihe für Vegetationskunde* **28**: 633–677.
- LEPŠÍ M, VÍT P, LEPŠÍ P, BOUBLÍK K & KOLAR F. 2009. *Sorbus portae-bohemicae* and *Sorbus albensis*, two new apomictic endemic species recognized based on revision of *Sorbus bohemica*. *Preslia* **81**: 63–89.
- LEPŠÍ M, VÍT P, LEPŠÍ P, BOUBLÍK K & SUDA J. 2008. *Sorbus milensis*, a new hybridogenous species from northwestern Bohemia. *Preslia* **80**: 229–244.
- LOHMEYER T. 1999. *Clitocybe truncicola* – ein holzbewohnender Trichterling aus der Sektion Candicantes. *Beiheft Zur Zeitschrift Für Mykologie* **9**: 69–72.
- MARTINCIĆ A. 2006. Moss flora of the Prokletije mountains (Serbia, Montenegro). *Hacquetia* **5**: 113–130.
- MAŠIĆ E, BARUDANOVIĆ S, ŽERO S, RAMIĆ E, MACANOVIC A, BOŠKAILO A & FEJZIĆ S. 2019. Diversity of diatoms in freshwater oligotrophic habitat types on Vranica Mountain (Bosnia and Herzegovina). *European Journal of Phycology* **54** (Suppl.): 128.

- NÉMETH C, BARABITS E & BÍLÁ J. 2016. New *Sorbus* subg. *Tormaria* (*S. latifolia* agg.) species from the southwestern part of the Transdanubian mountain range (Keszthely Mts, Hungary). *Studia Botanica Hungarica* **47**(2): 297–318.
- NIKETIĆ M. 2018. The first record of *Ephedra distachya* L. (Ephedraceae, Gnetophyta) in Serbia. Biogeography, coenology, and conservation. *Botanica Serbica* **42**(1): 123–138.
- PAPP B, ERZBERGER P, ÓDOR P, HOCK Z, SZÖVÉNYI P, SZURDOKI E & TÓTH Z. 2010. Updated checklist and red list of Hungarian bryophytes. *Studia Botanica Hungarica* **41**: 31–59.
- RANDJELOVIĆ V. 1999. *Typha shuttleworthii* Koch & Sonder. In: STEVANOVIC V (ed.), *Crvena knjiga flore Srbije* **1**, pp. 384–386, Ministarstvo za životnu sredinu Republike Srbije, Biološki fakultet Univerziteta u Beogradu, Zavod za zaštitu prirode Republike Srbije, Beograd.
- RANKOU H. 2011. *Epipactis purpurata*. The IUCN Red List of Threatened Species 2011. Published on the Internet <http://dx.doi.org/10.2305/IUCN.UK.2011-2.RLTS.T175965A7156460.en>. [Accessed 27 March 2020].
- RICH TCG, HOUSTON L, ROBERTSON A & PROCTOR MCF. 2010. *Whitebeams, Rowans and Service Trees of Britain and Ireland. A monograph of British and Irish Sorbus L. B.S.B.I. Handbook No. 14*. Botanical Society of the British Isles, London.
- ROBERTSON A, RICH TCG, ALLEN AM, HOUSTON L, ROBERTS C, BRIDLE JR, HARRIS SA & HISCKOK SJ. 2010. Hybridization and polyploidy as drivers of continuing evolution and speciation in *Sorbus*. *Molecular Ecology* **19**: 1675–1690.
- SABOVLJEVIĆ M & GRDOVIĆ S. 2009. Bryophyte diversity within urban areas: case study of the city of Belgrade (Serbia). *International Journal of Botany* **5**(1): 85–92.
- SOŠKA, T. 1949. Pregled mahovina i lišajeva u okolini Beograda. *Glasnik Prirodno-začinskog Muzeja Srpske Zemlje* **1–2**: 93–112.
- ŠPANIEL S, MARHOLD K & ZOZOMOVÁ-LIHOVÁ J. 2017. The polyploid *Alyssum montanum* - *A. repens* complex in the Balkans: a hotspot of species and genetic diversity. *Plant Systematics and Evolution* **303**: 1443–1465.
- ŠTEFĂNUȚ S. 2000. *Haplomitrium hookeri*, new species of Romania. *Acta Horti Botanici Bucurestiensis* **28**: 139–141.
- ŠTEFĂNUȚ S. 2007. Reports 8–11. In: NATCHEVA R (ed.), New bryophyte records in the Balkans: 1. *Phytologia Balcanica* **13**: 101–106.
- ŠTEFĂNUȚ S. 2010. Reports 18–28. In: NATCHEVA R (ed.), New bryophyte records in the Balkans: 6. *Phytologia Balcanica* **16**: 137–141.
- ŠTEFĂNUȚ S & GOIA I. 2012. Checklist and Red List of bryophytes of Romania. *Nova Hedwigia* **95**: 59–104.
- STOYNEVA-GÄRTNER MP, ISHEVA T, UZUNOV B & DIMITROVA P. 2015. Red list of Bulgarian algae. II. Microalgae. *Annuaire de l'Université de Sofia "St Kimment Ohridski", Botanique* **100**: 1–55.
- TEOFILOVSKI A. 2017. Contribution to knowledge of the flora of the Republic of Macedonia. *Botanica Serbica* **41**(1): 99–103.
- TOMOVIĆ G, ZLATKOVIĆ B, NIKETIĆ M, PERIĆ R, LAZAREVIĆ P, DURAKI Š, STANKOVIĆ M, LAKUŠIĆ D, ANAČKOV G, KNEŽEVIĆ J, SZABADOS K, KRIVOŠEJ Z, PRODANOVIĆ D, VUKOJIČIĆ S, STOJANOVIĆ V, LAZAREVIĆ M & STEVANOVIC V. 2009. Threat status revision of some taxa from "The Red Data Book of Flora of Serbia 1". *Botanica Serbica* **33**(1): 33–43.
- VUČKOVIĆ M. 1991. *Livadska i livadsko-stepska vegetacija Vršačkih planina*. Matica Srpska, Novi Sad.
- VUKIĆEVIĆ E. 1976. Rod. *Stipa* L. In: JOSIFOVIĆ M (ed.), *Flora Srbije* **8**, pp. 316–322, Srpska akademija nauka i umetnosti, Beograd.
- WRABER T. 1990. De *Gentiana nopscae* Jávorka notula. *Bioški vestnik* **38**(3): 39–46.
- ZIELIŃSKI J & VLADIMIROV V. 2013. *Sorbus ×latifolia* s.l. (Rosaceae) in the Balkan Peninsula and SW Asia. *Phytologia Balcanica* **19**(1): 39–46.
- ZLATKOVIĆ B & BOGOSAVLJEVIĆ S. 2014. Report on the new floristic data from Serbia. *Biologica Nyssana* **5**(2): 123–129.
- ZLATKOVIĆ B, RANDJELOVIĆ V & JUŠKOVIĆ M. 2007. Reports 136–138. In: VLADIMIROV V, DANE F, STEVANOVIC V & TAN K (ed.), New floristic records in the Balkans: 6. *Phytologia Balcanica* **13**(3): 433–455.

REZIME



Novi i značajni podaci o biljkama, algama i gljivama iz JI Evrope i susednih regiona, 2

Gordana TOMOVIĆ, Marko S. SABOVLJEVIĆ, Ivana ĐOKIĆ, Predrag PETROVIĆ, Vladan ĐORDJEVIĆ, Predrag LAZAREVIĆ, Ermin MAŠIĆ, Senka BARUDANOVIC, Sorin ŠTEFĀNUȚ, Marjan NIKETIĆ, Branislava BUTORAC, Jovana PANTOVIĆ, Alma HAJRUDINOVIĆ-BOGUNIĆ, Faruk BOGUNIĆ, Eva KABAŠ, Snežana VUKOJIČIĆ, Nevena KUZMANOVIĆ, Sanja Z. ĐJUROVIĆ i Uroš BUZUROVIĆ

Prikazani su novi i značajni podaci sa područja JI Evrope i susednih regiona o dijatomejskoj algi *Eunotia boreoalpina*; saprofitskoj gljivi *Clitocybe truncicola*; jetrenjači *Haplomitrium hookeri*, mahovini *Leptodon smithii*: monokotilama *Epipactis purpurata*, *Stipa tirsia*, *Typha laxmannii* i *T. shuttleworthii*; i dikotilama *Krascheninnikovia ceratoides*, *Polygonum albanicum* i *Sorbus latifolia*.

KLJUČNE REČI: novi prilog, *Clitocybe truncicola*, *Eunotia boreoalpina*, *Epipactis purpurata*, *Haplomitrium hookeri*, *Krascheninnikovia ceratoides*, *Leptodon smithii*, *Polygonum albanicum*, *Sorbus latifolia*, *Stipa tirsia*, *Typha laxmannii*, *Typha shuttleworthii*