


Klub mladih hemičara Srbije  
Serbian Young Chemists' Club

Srpsko hemijsko društvo  
Serbian Chemical Society



Šesta konferencija  
mladih hemičara Srbije

# ***Kratki izvodi radova***

Sixth Conference  
of the Young Chemists of Serbia  
***Book of Abstracts***

**Beograd, 27. oktobar 2018.**  
**Belgrade, 27<sup>th</sup> October 2018**



CIP - Каталогizacija u publikaciji  
Narodna biblioteka Srbije, Beograd

54(048)(0.034.2)  
577.1(048)(0.034.2)  
60(048)(0.034.2)  
66.017/.018(048)(0.034.2)

KONFERENCIJA mladih hemičara Srbije (6 ; 2018 ; Beograd)

Kratki izvodi radova [Elektronski izvor] = Book of Abstracts / Šestakonferencija mladih hemičara Srbije, Beograd, 27. oktobar 2018. = Sixth Conference of Young Chemists of Serbia, Belgrade, 27th October 2018 ; [organizatori Klub mladih hemičara Srbije [i] Srpsko hemijsko društvo =organizers Serbian Young Chemists' Club [and] Serbian Chemical Society] ; [urednici, editors Tamara Todorović ... et al.]. - Beograd : Srpskohemijsko društvo = Serbian Chemical Society, 2018 (Beograd :Razvojno-istraživački centar grafičkog inženjerstva TMF = Belgrade :Development and Research Centre of Graphic Engineering Faculty of Technology and Metallurgy). - 1 elektronski optički disk (CD-ROM) : tekst ; 12 cm

Sistemski zahtevi: Nisu navedeni. - Apstrakti na engl. jeziku. - Nasl. sa naslovne strane dokumenta. - Tiraž 120. - Bibliografija uz većinu apstrakata. - Registar.

ISBN 978-86-7132-072-6

1. Klub mladih hemičara Srbije (Beograd) 2. Srpsko hemijsko društvo (Beograd)

a) Хемија - Апстракти b) Биохемија - Апстракти c) Биотехнологија - Апстракти d) Наука о материјалима - Апстракти  
COBISS.SR-ID 269395724

## **6. KONFERENCIJA MLADIH HEMIČARA SRBIJE**

6<sup>th</sup> CONFERENCE OF THE YOUNG CHEMISTS OF SERBIA

**Beograd, 27. oktobar 2018. / BELGRADE, 27<sup>th</sup> OCTOBER 2018**

**KRATKI IZVODI RADOVA / BOOK OF ABSTRACTS**

*Izdaje / Published by*

**Srpsko hemijsko društvo / Serbian Chemical Society**

Karnegijeva 4/III, 11000 Beograd, Srbija

Tel./fax: +381 11 3370 467; www.shd.org.rs; office@shd.org.rs

*Za izdavača / For Publisher*

**Vesna MIŠKOVIĆ-STANKOVIĆ, predsednica Društva / President**

*Urednici / Editors*

**Tamara TODOROVIĆ**

**Ljubodrag VUJISIĆ**

**Jelena RADIVOJEVIĆ**

**Vuk FILIPOVIĆ**

*Dizajn, slog i kompjuterska obrada / Page Layout and Design*

**Vuk FILIPOVIĆ**

*Tiraž / Circulation*

**120 primeraka / 120 copies**

**ISBN 978-86-7132-072-6**

*Umnožavanje / Copying*

**Razvojno-istraživački centar grafičkog inženjerstva,**

**Tehnološko-metalurški fakultet, Karnegijeva 4, Beograd, Srbija**

Development and Research Centre of Graphic Engineering

Faculty of Technology and Metallurgy, Karnegijeva 4, Belgrade, Serbia

HA04 PE 2

## Optimization of solid-phase extraction for analysis of artificial sweeteners in water

Eleonora Gvozdić<sup>1</sup>, Ivana Matić Bujagić<sup>2</sup>, Ljiljana Tolić Stojadinović<sup>1</sup>, Svetlana Grujić<sup>2</sup>

<sup>1</sup>University of Belgrade, Innovation Centre of the Faculty of Technology and Metallurgy, Karnegijeva 4, 11000 Belgrade

<sup>2</sup>University of Belgrade, Faculty of Technology and Metallurgy, Karnegijeva 4, 11000 Belgrade

After decades of using artificial sweeteners in the food and pharmaceutical industry, their ubiquitous presence in the environment has been determined [1,2]. These substances are recognized as emerging pollutants due to limited data on their occurrence, environmental fate and ecotoxicity. Prior to analysis of artificial sweeteners at trace levels in water samples, it is necessary to isolate and concentrate analytes and clean-up the obtained extracts. In this paper, the optimization of the preparation method for the determination of saccharin, sucralose and neotame in water by solid-phase extraction (SPE) was performed. The extracts were analyzed by liquid chromatography–tandem mass spectrometry (LC–MS/MS). In the SPE optimization, the following parameters were tested: the sorbent type, the pH value and the volume of water sample and the elution solvent. The suitability of anion-exchange, polymeric, C18 and carbon sorbents for the extraction of selected sweeteners was assessed. The results showed that the highest recoveries were obtained using polymeric Oasis HLB cartridges, with copolymer poly(divinylbenzene-co-N-vinylpyrrolidone). Furthermore, it was determined that the pH value of water sample was crucial for extraction efficiency. For all three artificial sweeteners, the optimal pH value of the sample was 3. Finally, the sample volume of 100 ml ensured a high preconcentration factor without the sorbent breakthrough, and methanol was selected as the elution solvent.

### References

1. P. Arbeláez, F. Borrull, E. Pocurull, R.M. Marcé, *J. Chromatogr. A* **2015**, 1393, 106.
2. J. Spolestra, N.D. Senger, S.L. Shiff, *J. Environ. Qual.* **2017**, 46(6), 1434.

### Acknowledgments

This work was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (project no. 172007).