

EMEC21

21st European Meeting on Environmental Chemistry
November 30 – December 3, 2021, Novi Sad, Serbia

www.emec21.rs



Association of Chemistry
and the Environment



Serbian Chemical Society



Matica Srpska

Scientific Committee

Jan Schwarzbauer, president

Organisational Committee


Branimir Jovančević, president

Executive Committee

Vladimir Beškoski, president



BOOK OF ABSTRACTS



21st European Meeting
on Environmental Chemistry

EMEC 21



21st European Meeting
on Environmental Chemistry

BOOK OF ABSTRACTS
EMEC 21

November 30 – December 3, 2021

Novi Sad, Serbia



Book of Abstracts
21st European Meeting on Environmental Chemistry

Publisher

Serbian Chemical Society
Karnegijeva 4/III, Belgrade, Republic of Serbia

For the publisher

Dušan Sladić
President of the Serbian Chemical Society

Editors

Ivana Ivančev-Tumbas
Vladimir P. Beškoski
Aleksandra Šajnović

Cover page photo

Branko Lučić

Design and prepress

Beoživković, Belgrade

Printed by

RIS Studio, Belgrade

Circulation

150

ISBN

978-86-7132-078-8

Year

2021

Anthropogenic and Climate Influence on Land Degradation

N. Antić^{1,*}, M. Stefanović², N. Mijatović³, M. Kašanin-Grubin¹, G. Veselinović¹, S. Stojadinović¹, B. Jovančičević⁴. (1) University of Belgrade-Institute of Chemistry, Technology and Metallurgy, Department of Chemistry, Njegoševa 12, 11000 Belgrade, Republic of Serbia, (2) Innovation Center of the Faculty of Technology and Metallurgy, Karnegijeva 4, Belgrade, Serbia, (3) Institute for Testing of Materials IMS, Vojvode Mišića Bl. 43, 11000 Belgrade, Serbia, (4) University of Belgrade - Faculty of Chemistry, Studentski trg 12-16, 11000 Belgrade, Serbia; *nevena.antic@ihm.bg.ac.rs.



Badlands are areas with scarce or completely absent vegetation formed in a wide range of lithologies in different climate conditions and exposed to a wide range of geomorphological processes [1]. Generally, rapid evolution governed by erosion processes is a consequence of complex mineralogical and physico-chemical sediment composition and climate conditions. Because of that, badlands are often described as natural field laboratories and, furthermore, badland material is suitable for laboratory experiments that can, in controlled conditions, provide insight of changes that occur in the field.

As indicated above, beside lithology, climate is one of the most significant factors in badlands forming. Since human activities have great impact on the environment and since climate changes present one of the biggest environmental pollution problems nowadays, in this research badland material was exposed to different conditions with the aim of monitoring changes caused by extreme climate conditions and acid ice.

Three samples from badlands in China organized in six sets were treated with ice (representing snow) and acid ice (frozen acid rain) during fifteen cycles, dried in the oven for three cycles and afterwards again treated

with ice and acid ice for additional five cycles. After each cycle samples were photographed, so that physical changes can be tracked, while leachate was collected and analyzed for monitoring changes in its volume, pH, electrical conductivity (EC) and cation concentration.

Beside slight oscillations in parameters through cycles of samples treated with acid ice, extreme changes in observed parameters were not noticed neither between samples, nor between treatments. Leachate EC were a bit higher in samples treated with ice, leachate volume was higher for samples treated with acid ice, while pH was similar in both cases. Cation concentrations are similar in the leachate of all tested samples. In most of cases, the highest concentrations were measured at the beginning of the experiment, during the first two cycles or during the first “ice” cycles after drying. This indicates the high cation concentrations originate from the sediment surface or washing along the crack that appeared after drying.

Physical changes that occurred through cycles implied that heat/drought is more aggressive agent of sediment decay. Decay caused by ice is slower, not as aggressive as drought, but not negligible, causing noticeable and significant cracks and fissures of fragments.

This experiment confirmed that drought has high impact on sediment weathering, but more importantly, pointed out the impact of ice and its thawing, opening new questions about climate impact on forming, erosion processes and evolution of badlands which need to be further examined.

Acknowledgements

The authors would like to thank the Ministry of Education, Science and Technological Development of Republic of Serbia (Grant No: 451-03-9/2021-14/200026) for financial support.

References

- [1] Martínez-Murillo JF, Nadal-Romero E. 2018. *Perspectives on Badland Studies in the Context of Global Change. In: Badlands Dynamics in a Context of Global Change.* Elsevier Inc. p. 1–25.