

 УНИВЕРЗИТЕТ У БАЊОЈ ЛУЦИ

 UNIVERSITY OF BANJA LUKA

 ТЕХНОЛОШКИ ФАКУЛТЕТ

 FACULTY OF TECHNOLOGY



PROCEEDINGS

OCTOBER 21-22, 2022

ACADEMY OF SCIENCES AND ARTS OF THE REPUBLIC OF SRPSKA, BANJA LUKA, REPUBLIC OF SRPSKA, B&H

INTERNATIONAL SCIENTIFIC CONFERENCE

OF CHEMISTS, TECHNOLOGISTS AND ENVIRONMENTALISTS OF REPUBLIC OF SRPSKA

XIV CONFERENCE OF CHEMISTS, TECHNOLOGISTS AND ENVIRONMENTALISTS OF REPUBLIC OF SRPSKA BOOK OF PROCEEDINGS

Publisher: University in Banjaluka, Faculty of Technology

Editorial board: Borislav Malinović, PhD, dean

Design and computer processing Pero Sailović, PhD MSc Marina Rakanović MSc Đorđe Vujčić

CIP - Каталогизација у публикацији Народна и универзитетска библиотека Републике Српске, Бања Лука
66(082) 661:663/664(082) 502(082)
CONFERENCE of Chemists, Technologists and Environmentalists of Republic of Srpska (14 ; 2023) [Book of proceedings] : international scientific conference / XIV Conference of Chemists, Technologists and Environmentalists of Republic of Srpska ; [editorial board Borislav Malinović] Banja Luka : University in Banjaluka, Faculty of Technology, 2023 ([S.l : s.n.]) 313 crp. ; 24 cm
Библиографија уз сваки рад.
ISBN 978-99938-54-98-2
COBISS.RS-ID 137637377

Organizing Committee:

PhD Pero Sailović, president, PhD Darko Bodroža, general secretary, M.Sc Maja Milijaš secretary, M.Sc Dajana Dragić, M.Sc Branka Ružičić, M.Sc Marina Rakanović, M.Sc Maja Katić, Sanda Pilipović, M.Sc Nebojša Gorgi, Biljana Vasić, Sanja Novaković, M.Sc Jovanka Kotur *Students*: Vladimir Ivković, Jovan Savić, Nevena Janjić, Bojana Milinković, Danijela Lazić

Scientific Committee:

Dr Božana Odžaković, president, University of Banja Luka, **B&H**, Dr Nada Štrbac, co-president, University of Belgrade, Serbia, Dr Borislav Malinović, University of Banja Luka, B&H, Dr Vlada Veljković, University of Niš, Serbia, Dr Todor Vasiljević, Victoria University Melbourne, Australia, Dr Sanja Mahović-Poljačak, University of Zagreb, Croatia, Dr Csaba Horvath, University Obuda, Budapest, Hungary, Dr Mihail Kochubovski, University of Skopje, Macedonia, Dr Massimiliano Fenice, Universityt Della Tuscia, Italy, Dr Georgij Petriaszwili, Warshav University of Technology, Poland, Dr Mira Vukcević, University of Monte Negro, Monte Negro, Dr Ondrej Panák, University of Pardubice, Czech Republic, Dr Pospiech Matej, University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic, , Dr Dani Dordevic, University of Veterinary and Pharmaceutical Sciences, Brno, Czech Republic, Dr Iskren Spiridonov, University of Chemical Technology and Metallurgy, Bulgaria, Dr Laura Benea, West University of Timisoara, Romania, Dr Savvas G. Vassiliadis, University of Piraeus, Greece, Dr Helena Prosen, University of Ljubljana, Slovenia, Dr Srecko Stopic, RWTH University Aachen, Germany, Dr Maria Iosune Cantalejo, UPNA, Spain, Dr Jurislav Babić, University of Osijek, Croatia, Dr Svetozar Milosavić, University of Kosovska Mitrovica, Serbia, Dr Petar Uskoković, University of Belgrade, Serbia, Dr Mitja Kolar, University of Ljubljana, Slovenia, Dr Dragiša Savić, , University of Niš, Serbia, Dr Dragan Vujadinović, University of East Sarajevo, **B&H**, Dr Biljana Pajin, University of Novi Sad, Serbia, Dr Sead Catić, University of Tuzla, B&H, Dr Husein Vilić, University of Bihać, B&H, Dr Sanjin Gutić, University of Sarajevo, B&H, Dr Goran Trbić, University of Banja Luka, B&H, Dr Milica Balaban, University of Banja Luka, B&H, Dr Ljiljana Vukić, University of Banja Luka, **B&H**, Dr Ljiljana Topalić-Trivunović, University of Banja Luka, **B&H**, Dr Slavica Sladojević, University of Banja Luka, **B&H**, Dr Pero Dugić, University of Banja Luka, **B&H**, Dr Zoran Kukrić, University of Banja Luka, **B&H**, Dr Slavica Grujić, University of Banja Luka, **B&H**, Dr Milorad Maksimović, University of Banja Luka, B&H, Dr Branka Rodić-Grabovac, University of Banja Luka, B&H, Dr Rada Petrović, University of Banja Luka, B&H, Dr Dragana Grujić, University of Banja Luka, B&H, Dr Svjetlana Janjić, University of Banja Luka, B&H, Dr Zora Levi, University of Banja Luka, **B&H**, Dr Ladislav Vasilišin, University of Banja Luka, **B&H**

NOTE:

The authors have full responsibility for the originality and content of thier own papers



UNIVERSITY OF BANJA LUKA TEXHONOШKИ ФАКУЛТЕТ FACULTY OF TECHNOLOGY



International scientific conference

"XIV CONFERENCE OF CHEMISTS, TECHNOLOGISTS AND ENVIRONMENTALISTS OF REPUBLIC OF SRPSKA"

under the auspices of



MINISTRY OF SCIENTIFIC AND TECHNOLOGICAL DEVELOPMENT, HIGHER EDUCATION AND INFORMATION SOCIETY

MINISTRY OF AGRICULTURE, FORESTRY AND WATER MANAGEMENT



ACADEMY OF SCIENCES AND ARTS OF THE REPUBLICA OF SRPSKA

SPONSORS

OPTIMA GROUP and OIL RAFINERY MODRIČA HOFSTETTER ENVIRONMENTAL SRL SINEX LABORATORY PREHTEH d.o.o. **DESTILACIJA** ad EAST CODE d.o.o. MUNICIPALITY OF ČELINAC J.P. DEP-OT ASSOCIATION OF TECHNOLOGY ENGINEERS OF REPUBLIC OF SRPSKA EM PLUS d.o.o. KESO GRADNJA d.o.o. MARKWAY d.o.o. EURO-LAB V d.o.o. EKO-EURO TIM d.o.o. MB-IMPEX d.o.o. TRGO FORTUNA PLUS d.o.o. EURO-INSPEKT d.o.o. ABC PROJEKT d.o.o ŠTAMPARIJA PETROGRAF **CSK PRINT**

CONTENTS

GENERAL AND APPLIED CHEMISTRY
SYNTHESIS AND ANTIPROLIFERATIVE ACTIVITY OF NITRO AND AMINO SUBSTITUTED BENZIMIDAZOLES AND BENZOTHIAZOLES
PHOTOLYSIS OF NADOLOL IN THE AQUATIC ENVIRONMENT: INSIGHTS INTO THE EFFECT OF BICARBONATES
CAN E-CIGARETTES BE A THREAT TO ENVIRONMENT: HEAVY METALS ANALYSIS IN E-CIGARETTES FILTERS
COMPARISON OF THREE EXTRACTION PROCEDURES FOR QUANTIFICATION OF Cu, Mn, Ni, Pb AND Zn IN THE PLANTS
ANTIPROLIFERATIVE AND ANTIBACTERIAL ACTIVITY IN VITRO OF AMIDINO SUBSTITUTED 2- NAPHTHYLBENZAZOLES
SOLVATOCHROMIC PROPERTIES OF NEWLY SYNTHESIZED ARYL AZO PYRIDONE DYES
CHEMICAL ENGINEERING
BIOBASED UNSATURATED POLYESTER RESINS REINFORCED WITH NATURAL FILLERS
POSSIBILITIES OF CROSSLINKING SILICONE MATERIALS IN EXCESS OF CROSSLINKER WITH TWO TYPES OF FILLERS
Darko Manjenčić, Marko Paić, Vesna Cvijetinović, Vladan Mićić, Anja Manjenčić, Duško Kostić, Pero Dugić
PHOSPHATE GLASSY FERTILIZERS
CHEMICAL TECHNOLOGY
MINEROLOGICAL AND CHEMICAL CHARACTERIZATION OF CLAY FROM THE ZAGONI SITE MUNICIPALITY OF BRATUNAC
THE PHOTOSTABILITY AND DYEING ABILITY OF SOME ANTHRAQUINONE REACTIVE DYES FOR COTTON AND PAPER
COMPARATIVE ANALYSIS OF BLEACHING OF SUNFLOWER OIL WITH COMMERCIAL BLEACHING EARTH AND BENTONITE POWDER ACTIVATED WITH SULFURIC ACID
BIOTECHNOLOGY
THE IMPACT OF pH VALUE AND TEMPERATURE ON POLYPHENOL RECOVERY, ANTIOXIDANT CAPACITY AND PHYSICAL PROPERTIES OF SERPYLLI HERBA'S WASTE EXTRACTS
THE POSSIBILITY OF ANTIBIOTIC AZALOMYCIN B PRODUCTION BY COMBINING BENZOYL HYDRAZONE DERIVATIVES AND WASTE RAPESEED OIL GLYCEROL

Jovan Ćirić, Slavica Ilić, Sandra Konstantinović, Đorđe Lazarević, Marko Živković, Nikola Stanković

FOOD TECHNOLOGY
RED WINE COLOR AND STABILITY AS FUNCTION OF SIZE AND CONTENT OF SILICA GEL PARTICLES110 Tanja Šaran, Dragan Vujadinović, Zoran Petković, Milan Vukić, Vesna Gojković, Vladimir Tomović
THE SEASONAL VARIATIONS OF BIOCHEMICAL COMPOSITION OF COW'S MILK ON THE SOUTHERN SERBIA TERRITORY
Jovan Ćirić, Slavica Ilić, Aleksandar Veličković, Ivan Stojković, Dragana Stanisavljević, Nebojša Milosavljević
THE EFFECTS OF ADDITION OF FLAXSEED TO THE CHARACTERISTICS OF BREAD122
Gordana Jovanović, Ana Vasić, Ljubica Mijić, Ana Matić, Bojan Damnjanović
HABITS OF FLUID INTAKE WITH STUDENTS
QUALITY CONTROL AND FOOD SAFETY134
THE CONTENT OF TOTAL IRON IN <i>Urtica dioica L.</i> FROM DIFFERENT LOCALITIES IN SERBIA
THE PRESENCE OF PHOSPHATE IN MEAT PRODUCTS FROM THE MARKET OF REPUBLIC OF SRPSKA 143 Biljana Pećanac, Dragan Brenjo
TEXTILE ENGINEERING
OPTIMIZATION OF THE CLASSICAL DYEING PROCESS OF PES KNITWEAR
INFLUENCE OF KNIT PATTERN ON DIMENSIONAL STABILITY OF KNITTED FABRICS
GRAPHIC ENGINEERING AND DESIGN166
A METHOD OF RANKING RESPONDENTS ACCORDING TO SENSE OF COLOR DIFFERENCES
AN OVERVIEW OF VARIOUS POSSIBILITIES OF 3D PRINTING TECHNIQUES IN THE PRODUCTION OF FLEXOGRAPHIC PRINTING PLATES
CONTEMPORARY APPLICATIONS OF PULP MOULDED PACKAGING
Gojko Vladić, Nemanja Kašiković, Živko Pavlović, Stefan Đurđević, Gordana Bošnjaković, Teodora Gvoka
INFLUENCE OF THE SUBSTRATE ON THE PERMANENCE OF PRINTING INK
REMARKS ABOUT PRINTING AUTOMATION
ENVIRONMENTAL ENGINEERING AND ECOLOGY
DEPOSITION OF GOETHITE AND FERRIHYDRITE ONTO EXPANDED VERMICULITE SURFACE: CHEMICAL PROPERTIES AND POTENTIAL APPLICATION
REMOVAL OF XENOBIOTICS FROM WASTEWATERS USING PHOTOLYSIS UNDER SUN-LIGHT IRRADIATION: EXPERIMENTAL APPROACH AND PROCESS DESIGN

Aleksandar Jovanović, Mladen Bugarčić, Nataša Knežević, Jovana Bošnjaković, Jelena Lukić, Antonije Onjia, Aleksandar Marinković

MATERIAL	SCIEN	CE AND	ME	FALLURGY	7	•••••	•••••	••••••	
SIMULATION	OF THE	IMPACT	OF P	REHEATING	TEMPERATURE	ON	RAILWAY	ALUMINOT	HERMIC

 Tamara T. Tadić, Bojana M. Marković, Mila V. Ilić, Aleksandra B. Nastasović, Antonije E. Onjia

Original scientific article

REMOVAL OF XENOBIOTICS FROM WASTEWATERS USING PHOTOLYSIS UNDER SUN-LIGHT IRRADIATION: EXPERIMENTAL APPROACH AND PROCESS DESIGN

Aleksandar Jovanović¹, Mladen Bugarčić¹, Nataša Knežević², Jovana Bošnjaković³, Jelena Lukić⁴, Antonije Onjia², Aleksandar Marinković²

¹Institute for Technology of Nuclear and Other Mineral Raw Materials, 11000 Belgrade, Serbia ²University of Belgrade, Faculty of Technology and Metallurgy, 11000 Belgrade, Serbia ³Research and Development Institute Lola L.T.D., 11030 Belgrade, Serbia ⁴Innovation Center of Faculty of Technology and Metallurgy, 11000 Belgrade, Serbia

Abstract

Conventional wastewater treatment processes are increasingly difficult to fulfill strict standards of achieving the required quality. Therefore, it is necessary to apply processes that in an efficient and economically justified way reach the necessary limits in order to discharge wastewater into the recipient. The process of photolysis represents the degradation of the starting pollutant practically to water and carbon dioxide. The whole process takes place under the action of a UV lamp that imitated solar irradiation. The process is environmentally justified since, unlike some other advanced oxidation processes - AOPs, there is no use of additional, hazardous, chemicals which possibly results in the precipitates formation that are leading to further treatment. In this paper, photolytic degradation of pesticide thiophanate methyl was performed in doubled wall quartz reactor equipped with a thermoregulation system. Reaction was forced under Osram ULTRA VITALUX UV lamp (300W). Kinetics of photodecomposition process was followed by pseudo-first order. In certain time periods, specimens were sampled, filtered and monitored by UV spectrophotometer Shimadzu 1800 with an aim to determine the concentration of xenobiotic. Complete degradation of present xenobiotics was obtained after 240 min. With an aim to enhance the degradation process, hydrogen peroxide was added as a boosting agent which cause a shortening in demanded time (150 min) for complete degradation. Verification of the success of decomposition was confirmed by the obtained values of chemical oxygen demand (COD), which reveal that the established system has a basis for real application in industrial or municipal wastewater.

Keywords: wastewater treatment, pesticide degradation, photolysis, AOPs, COD.

Introduction

Due to the excessive or improper use of pesticides, the rising demand for food production worldwide has not only produced a major decline in food quality and severe environmental effects (Abhilash & Singh, 2009). Various pesticides used in agricultural techniques to assure crop quality and food safety might cause the release of these compounds into non-target environmental matrices, such as freshwater bodies (Bustos et al, 2019). This is a serious issue because pesticides are typically recognized to have harmful qualities that can make exposure to these chemicals potentially lethal. In terms of human health, pesticide exposure has been linked to an increase in the occurrence of some forms of cancer and birth abnormalities, among other disorders (Kaur & Kaur, 2018). Furthermore, it is generally understood that these substances might pose a health danger to other living species (Damalas & Eleftherohorinos, 2011). A fungicide called thiophanate methyl was frequently applied

to crops to prevent fungal infections (Correia et al., 2016). It is not particularly persistent in soil or water systems, has a low aqueous solubility, and is not particularly volatile. Although it is not poisonous to mammals, it can irritate, cause skin sensitivity, and even be mutagenic (Jia et al., 2020). Most aquatic species and earthworms are poisonous to it, while birds and bees are less susceptible. Due to high water pollution with different organic compounds, many researchers are trying their best to develop techniques for the removal and degradation of them. Wastewaters from industrial factories and agricultural fields present a tremendous problem for modern society, in the area of health care and sustainability (Ungureanu et al., 2020). Therefore, oxidative process like photolysis is increasingly used in smaller wastewater treatment plants (Liu et al., 2011; Anisuzzaman et al., 2022). This process provides the degradation of many classes of organic compounds and biological pollutants. Photolysis is a process of degradation of various pollutants under the action of UV irradiation, where the parent molecule is converted into smaller ones.

The goal of this study was to thoroughly investigate the degradation of thiophanate methyl (TPM) under simulated sunlight in order to develop an appropriate technique for its removal from water. Measurement of chemical oxygen demand (COD) will help in further explanations of process efficiency. The effects of hydrogen peroxide addition and the distance of the UV lamp on degradation were examined. The reaction kinetics was further investigated using UV-VIS.

Materials and Methods

All used chemicals are analytically pure and without the need for further purification. Pesticide thiophanate methyl was obtained from Sigma Aldrich (European Commission, 2022). Hydrogen peroxide was also acquired from Sigma Aldrich. Photolytic experiments were performed in a doubled-wall quartz glass reactor (Figure 1). For irradiation of pesticide solutions, Osram Ultra Vitalux UV lamp nominal wattage of 300 W (UVA : UVB = 13.6 : 3) was used. During irradiation, the reactor is consistently cooled down by coolant water in the reactor jacket to avoid uncontrolled heating of the treated solutions. Photolytic tests in this study were carried out with starting TPM concentrations of 5 and 10 mg dm⁻³. For initial experiments, in the reactor 150 dm⁻³ of the abovementioned fungicide solutions were added. In certain periods, 3.0 dm⁻³ of the solution was taken and recorded via UV spectrophotometry. All measurements were taken in triplicate. In photolytic experiments, the concentration of TPM was measured using UV-VIS Spectrophotometer 1800, Shimadzu, Japan. COD was measured by EPA 410.4: 1993 standard.




Figure 1. The reactor used for photolytic experiments

Results and discussion

At low concentrations of the organic pollutant, a pseudo-first order equation can be used to estimate the kinetics of photolytic reactions (Yusuff et al., 2020). In the Figure 2 are shown obtained curves with different symbols for all tests. They are representing the degradation of the present pollutant by sun-imitated irradiation and by sun-imitated irradiation with an addition 5% solution of H_2O_2 .




Figure 2. Degradation of TPM under different conditions

Photolytic tests reveal the relatively average degradation of TPM, which indicates the stability of this pesticide (which is in agreement with literature from Introduction). Results show that for both TPM concentrations, 5 and 10 mg dm⁻³, the needed time for complete decomposition was 240 min.

This is the reason for advancing photolytic reaction with the addition of H_2O_2 in order to fasten degradation of presented organic. The formation of peroxide radicals reduces the duration of the process due to accelerated oxidation of the pollution, which drives the degradation reaction and justify the necessity for an advanced process. Therefore, acquired results prove usage of this combination -150 min as the required time for total decomposition of TPM (Figure 2).

By varying the distance of the lamp from reactor (from 100 to 400 mm), it was concluded that a distance of 200 mm has the best impact on degradation reactions. When the distance was the higher then 200 nm, photolytic degradation was lower by approximately 15% for 300 mm and approximately 25% for 400 mm. This can be explained by dissipation of emitted light. Contrastingly, when the gap between the rector and UV lamp was smaller, solution of TPM started to evaporate because of inability of the system to cool the working solution.

Measurement of COD values are shown in Figure 3.




Figure 3. Monitoring of COD values prior and after the degradation of TPM

Determined initial value of COD for TPM solution was 1255 mg dm⁻³. Following the 240 min of photolytic reactions, obtained values were 510 and 420 for 10 and 5 mg dm⁻³, respectively. After the addition of 5% solution of H₂O₂, values of COD were decreased, 120 and 80 for 10 and 5 mg dm⁻³, respectively. This shows that the used system showed significant improvements.

Conclusion

Typical wastewater treatment systems are becoming increasingly difficult to meet stringent quality standards. As a result, in order to discharge wastewater into the recipient, processes that are efficient and economically justified must be used. Therefore, entire procedure is carried out under the influence of a UV light that simulates sun irradiation. The kinetics of the photodecomposition process was followed by pseudo-first order. Complete degradation of the present TPM was obtained after 240 min. With an aim to enhance the degradation process, hydrogen peroxide was added as a boosting agent which cause a shortening in demanded time (150 min) for complete degradation. Also, it was concluded that the optimal reactor-to-lamp distance was 200 mm. The estimated chemical oxygen demand (COD) values were used to evaluate the decomposition's success: 80 mg dm⁻³ for the concentration of TPM of 5 mg dm⁻³, which reveal that the established system has a basis for real application in industrial or municipal wastewater.

References

- Abhilash, P. C. & Singh, N. (2009). Pesticide use and application: an Indian scenario. *Journal of Hazardous Materials*, *165*(1–3), 1–12. https://doi.org/10.1016/j.jhazmat.2008.10.061
- Anisuzzaman, S. M., Joseph, C. G., Pang, C. K., Affandi, N. A., Maruja, S. N. & Vijayan, V. (2022). Current Trends in the Utilization of Photolysis and Photocatalysis Treatment Processes for the Remediation of Dye Wastewater: A Short Review. *ChemEngineering*, 6(4). https://doi.org/10.3390/chemengineering6040058
- Bustos, N., Cruz-Alcalde, A., Iriel, A., Fernández Cirelli, A. & Sans, C. (2019). Sunlight and UVC-254 irradiation induced photodegradation of organophosphorus pesticide dichlorvos in aqueous matrices. *Science of the Total Environment*, 649, 592–600.

https://doi.org/10.1016/j.scitotenv.2018.08.254

- Correia, M., Rodrigues, M., Paíga, P. & Delerue-Matos, C. (2016). Fungicides. In B. Caballero, P. M. Finglas, & F. B. T.-E. of F. and H. Toldrá (Eds.) (pp. 169–176). Oxford: Academic Press. https://doi.org/https://doi.org/10.1016/B978-0-12-384947-2.00342-1
- Damalas, C. A. & Eleftherohorinos, I. G. (2011). Pesticide exposure, safety issues, and risk assessment indicators. *International Journal of Environmental Research and Public Health*, 8(5), 1402–1419. https://doi.org/10.3390/ijerph8051402

European Commission. (2022). SDS Thiophanate-methyl.

- Jia, K., Cheng, B., Huang, L., Xiao, J., Bai, Z., Liao, X. & Lu, H. (2020). Thiophanate-methyl induces severe hepatotoxicity in zebrafish. *Chemosphere*, 248(28), 125941. https://doi.org/10.1016/j.chemosphere.2020.125941
- Kaur, K. & Kaur, R. (2018). Occupational Pesticide Exposure, Impaired DNA Repair, and Diseases. Indian Journal of Occupational and Environmental Medicine, 22(2), 74–81. https://doi.org/10.4103/ijoem.IJOEM_45_18
- Liu, Y. S., Ying, G. G., Shareef, A. & Kookana, R. S. (2011). Photolysis of benzotriazole and formation of its polymerised photoproducts in aqueous solutions under UV irradiation. *Environmental Chemistry*, 8(2), 174–181. https://doi.org/10.1071/EN10141
- Ungureanu, N., Vlăduț, V. & Voicu, G. (2020). Water Scarcity and Wastewater Reuse in Crop Irrigation. *Sustainability*, *12*(21), 9055. https://doi.org/10.3390/su12219055
- Yusuff, A. S., Taofeek Popoola, L. & Aderibigbe, E. I. (2020). Solar photocatalytic degradation of organic pollutants in textile industry wastewater by ZnO/pumice composite photocatalyst. *Journal of Environmental Chemical Engineering*, 8(4), 103907. https://doi.org/https://doi.org/10.1016/j.jece.2020.103907