

EMEC21

21st European Meeting on Environmental Chemistry
November 30 – December 3, 2021, Novi Sad, Serbia

www.emec21.rs



Association of Chemistry
and the Environment



Serbian Chemical Society



Matica Srpska

Scientific Committee

Jan Schwarzbauer, president

Organisational Committee


Branimir Jovančičević, president

Executive Committee

Vladimir Beškoski, president



BOOK OF ABSTRACTS



21st European Meeting
on Environmental Chemistry

EMEC 21



21st European Meeting
on Environmental Chemistry

BOOK OF ABSTRACTS
EMEC 21

November 30 – December 3, 2021

Novi Sad, Serbia



Book of Abstracts
21st European Meeting on Environmental Chemistry

Publisher

Serbian Chemical Society
Karnegijeva 4/III, Belgrade, Republic of Serbia

For the publisher

Dušan Sladić
President of the Serbian Chemical Society

Editors

Ivana Ivančev-Tumbas
Vladimir P. Beškoski
Aleksandra Šajnović

Cover page photo

Branko Lučić

Design and prepress

Beoživković, Belgrade

Printed by

RIS Studio, Belgrade

Circulation

150

ISBN

978-86-7132-078-8

Year

2021

The Response of Badland Materials from Spain with Different Mineralogical Content on Seasonal Changes

M. Stefanović^{1,*}, B. Jovančičević², A. Šajnović³, F. Gallart⁴, M. Moreno-de las Heras⁴, N. Antić³, M. Kašanin-Grubin³. (1) University of Belgrade, Innovation Center of the Faculty of Technology and Metallurgy, Karnegijeva 4, Belgrade, Serbia, (2) University of Belgrade, Faculty of Chemistry, Studentski trg 12-16, Belgrade, Serbia, (3) University of Belgrade, Institute of Chemistry, Technology and Metallurgy, Njegoseva 12, Belgrade, Serbia, (4) Institute of Environmental Assessment and Water Research, Carrer de Jordi Girona 18-26, Barcelona, Spain, *mstefanovic@tmf.bg.ac.rs.



Fig.1. Samples after finished experiments.

Badlands are areas with limited vegetation, reduced or no human activity, and a great variety of geomorphic processes [1]. Badland materials have a different response to the same environmental conditions, because of differences in their mineralogical and physico-chemical characteristics. Many studies show that smectite-poor sediments are more resistant to different weathering treatments of freezing, thawing, wetting, and drying, than smectite-rich materials [2,3]. In this paper, three unweathered samples of badlands from Spain were analyzed with the aim of monitoring, but also comparing physico-chemical changes caused by simulations of changes in climatic conditions. Selected sediment samples had different compositions. Besides quartz and calcite, the first sample is composed of smectite and gypsum (3 UW), the second of smectite (4 UW), while the third sample is composed of neither smectite nor gypsum (5 UW). The experiment setup was designed in the way that each sample had three sub-samples, a sample for simulation of rain, snow, and a control sample (Figure 1). Sample_rain was treated with a rain intensity of ~850 ml/h for 10 minutes (~140 ml), while sample_snow was treated with crushed ice (~150 g). After precipitation simulations snow were put samples were placed in a climate chamber at -3 °C together with a control sample. This was

repeated for 15 cycles. Every cycle was documented with photographs. The leached solution was collected and its volume, pH, electrical conductivity (EC), and ion concentrations were measured. The second part of the experiment was based on exposing the samples after wetting to higher temperatures, 50 °C. It was done in 8 cycles. FESEM and BET analyzes were performed for each sample before and after the experiments. The 3 UW samples had significantly different leachate pH and EC, while the leachate volume was similar for all samples during the experiment. Sulphate ions were leached in the highest concentrations during the whole experiment from the sample with both smectite and gypsum present. The sample with smectite has shown the highest disintegration of the structure, especially after the simulation of snow. The sample with smectite and gypsum has shown a lower degree of degradation than sample 3 UW due to the content of gypsum which increases the weathering resistance of the material. Sample 5 UW has shown the lowest degradation of the structure along with the weathering cycles. This study has proven that both mineralogical and physico-chemical properties of sediments are important for predicting their response to variable climate factors.

Acknowledgements

This work was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract No. 451-03-9/2021-14/200287).

References

- [1] J. F. Martínez-Murillo, E. Nadal-Romero, Perspectives on Badland Studies in the Context of Global Change in Badland Dynamics in the Context of Global Change, E. Nadal-Romero, J. F. Murillo, N. Kuhn, Elsevier, 2018, 1
- [2] M. Kasanin-Grubin, M. Della Seta, F. Troini, F. Versagi, 2018. The role of lithology and soils: parent material controls to badland development in Badland Dynamics in the Context of Global Change, E. Nadal-Romero, J. F. Murillo, N. Kuhn, Elsevier, 2018, 61
- [3] M. Kasanin-Grubin, *Catena*, 106 (2013) 54.